

THE EIGEN-PROBLEM FOR SOME SPECIAL NEAR-TOEPLITZ CENTRO-SKEW TRIDIAGONAL MATRICES

by

Antonio Behn

Department of Mathematics
University of Chile

Kenneth R. Driessel

Mathematics Department
Iowa State University

Irvin R. Hentzel

Mathematics Department
Iowa State University

ABSTRACT. Let $n \geq 2$ be an integer. Let R_n denote the $n \times n$ tridiagonal matrix with -1 's on the sub-diagonal, 1 's on the super-diagonal, -1 in the $(1, 1)$ entry, 1 in the (n, n) entry and zeros elsewhere. We find the eigen-pairs of the matrices R_n .

TABLE OF CONTENTS

- Introduction
- A Reduction
- Tridiagonal Toeplitz Matrices
- The Eigenvalues
- Acknowledgements
- References

Date: February 1, 2011.

1991 *Mathematics Subject Classification.* Primary: 15B05 Toeplitz, Cauchy, and related matrices; Secondary: 15B35 Sign pattern matrices; 15A18 Eigenvalues, singular values, and eigenvectors.

Key words and phrases. Tridiagonal, Toeplitz, eigenvalue, eigenvector, centro symmetric, centro skew symmetric, sign pattern.

INTRODUCTION

We consider some special n -by- n , near-Toeplitz, tridiagonal matrices with entries from the set $\{0, 1, -1\}$. In particular, we consider tridiagonal matrices having the following form:

$$R_n := \text{Tridiag}(\text{subdiag}, \text{diag}, \text{supdiag})$$

where

- $\text{subdiag} := (-1, -1, \dots, -1)$,
- $\text{diag} := (-1, 0, 0, \dots, 0, 0, 1)$, and
- $\text{supdiag} := (1, 1, \dots, 1, 1)$.

In other words, R_n is the tridiagonal matrix that has all -1 's on the subdiagonal, all 0 's on the diagonal except for a -1 in the $(1,1)$ entry and a 1 in the (n,n) entry, and has all 1 's on the superdiagonal. For example, when $n := 4$ we have:

$$R_4 := \begin{pmatrix} -1 & 1 & 0 & 0 \\ -1 & 0 & 1 & 0 \\ 0 & -1 & 0 & 1 \\ 0 & 0 & -1 & 1 \end{pmatrix}.$$

We find the eigenvalue-vector pairs (or **eigen-pairs** for short) of the matrices R_n . In particular, we prove the following result:

Proposition 1. *The eigenvalues of the matrix R_n are 0 and $2i \cos(j\theta)$ for $j = 1, \dots, n-1$ where $\theta := \pi/n$.*

Remark: If n is even then, 0 is an eigenvalue with multiplicity 2 .

Let $\mathcal{T}_n$ denote the set of $n \times n$ tridiagonal real matrices that satisfy the following conditions: the sub-diagonal is negative, the super-diagonal is positive, the $(1,1)$ entry is negative, the (n,n) entry is positive and all other entries are zero. Drew, et al(2000) conjectured that this sign pattern class contains matrices with arbitrary spectra. Note that the matrix R_n is in this class. They provided evidence for the conjecture. (Elsner, et al(2003) provided further evidence.) We believe that understanding the properties of R_n may be an important step toward understanding the arbitrary spectrum conjecture for the sign pattern class $\mathcal{T}_n$.

Here is a summary of the contents. In the section with title ‘‘A Reduction’’, we show that R_n is similar to a near skew-symmetric, tridiagonal, Toeplitz matrix. In the section with title, ‘‘Tridiagonal Toeplitz Matrices’’ we review the solution of the eigen-problem for such matrices; in particular, we determine the eigen-pairs for a skew-symmetric

tridiagonal Toeplitz matrix. In the section with title “The Eigenvalues”, we determine the eigenvalues of R_n ; in other words, we prove the proposition given above.

Aside: Let E be the $n \times n$ matrix defined by $E(i, j) := \delta(i + j, n + 1)$, for $1 \leq i, j \leq n$, where δ is the Kronecker delta. This matrix is called the **exchange matrix** or the **flip matrix**.

Let P be an $n \times n$ matrix. Then P is a **centro-symmetric matrix** if $EPE = P$ and P is a **centro-skew matrix** if $EPE = -P$.

Here is an older reference on centro-symmetry: Weaver(1985). Here is a more recent reference: Trench(2004). These papers contain further references. We shall not use the theory of centro-symmetric matrices in this paper. *EndAside.*

A REDUCTION

Recall that we are considering special matrices R_n where R_n is the tridiagonal matrix which has all -1 's on the subdiagonal, has all 0 's on the diagonal except for a -1 in the $(1,1)$ entry and a 1 in the (n,n) entry, and has all 1 's on the superdiagonal. We want to find the eigenvalues and eigenvectors of these matrices.

Here is another description of R_n . Let Z_n denote the **lower shift matrix** which is the matrix that has 1 's on the subdiagonal and 0 's elsewhere. For example, when $n = 4$, we have

$$Z_4 := \begin{pmatrix} 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}.$$

Aside: Here we follow notation and terminology used by T. Kailath. See, for example, Kailath, Kung and Morf(1979a), Kailath, Kung and Morf(1979b) or Kailath and Sayed(1999). *EndAside.*

Note that

$$R_n = Z_n^T - Z_n - e_1 e_1^T + e_n e_n^T$$

where e_k denotes the column vector which has 1 in the k th coordinate and 0 's elsewhere.

Proposition 2. Reduction. *Let $S_n := I_n + Z_n$ where I_n is the $n \times n$ identity matrix. Then*

$$S_n^{-1} R_n S_n = K_n + e_n e_{n-1}^T$$

where $K_n := Z_n^T - Z_n$.

Remark: Here is a picture of S_4 :

$$S_4 := \begin{pmatrix} 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \end{pmatrix}$$

Note that $I_n + Z_n$ is invertible since Z_n is nilpotent. In fact

$$(I_n + Z_n)^{-1} = I_n - Z_n + Z_n^2 - \cdots \pm Z_n^{n-1}.$$

When $n = 4$, for example, the conclusion of the proposition is

$$\begin{aligned} & \begin{pmatrix} 1 & 0 & 0 & 0 \\ -1 & 1 & 0 & 0 \\ 1 & -1 & 1 & 0 \\ -1 & 1 & -1 & 1 \end{pmatrix} \begin{pmatrix} -1 & 1 & 0 & 0 \\ -1 & 0 & 1 & 0 \\ 0 & -1 & 0 & 1 \\ 0 & 0 & -1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \end{pmatrix} \\ &= \begin{pmatrix} 0 & 1 & 0 & 0 \\ -1 & 0 & 1 & 0 \\ 0 & -1 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{pmatrix}. \end{aligned}$$

Proof. In order to reduce the number of symbols in this proof, we omit the subscript n on I , Z , R and S .

Note that the equation in the conclusion of the proposition is equivalent to

$$(*) \quad RS = SK + Se_n e_{n-1}^T.$$

Also note

$$\begin{aligned} RS &= (K - e_1 e_1^T + e_n e_n^T)S \\ &= KS + (e_n e_n^T - e_1 e_1^T)S \\ &= SK + [K, S] + (e_n e_n^T - e_1 e_1^T)S \end{aligned}$$

where $[X, Y] := XY - YX$.

Hence, to prove (*), we need only prove the following

$$\textit{Claim: } [K, S] = Se_n e_{n-1}^T + (e_1 e_1^T - e_n e_n^T)S$$

We have

$$\begin{aligned} [K, S] &= [K, I + Z] = [K, Z] = [Z^T - Z, Z] = [Z^T, Z] \\ &= Z^T Z - Z Z^T = \text{Diag}(1, 1, \dots, 1, 0) - \text{Diag}(0, 1, \dots, 1, 1) \\ &= e_1 e_1^T - e_n e_n^T. \end{aligned}$$

We also have

$$\begin{aligned}
 & S e_n e_{n-1}^T + (e_1 e_1^T - e_n e_n^T) S \\
 &= (I + Z) e_n e_{n-1}^T + (e_1 e_1^T - e_n e_n^T) (I + Z) \\
 &= e_1 e_1^T - e_n e_n^T.
 \end{aligned}$$

□

TRIDIAGONAL TOEPLITZ MATRICES

In this section, we consider the skew symmetric tridiagonal matrices $K_n := Z_n^T - Z_n$. We determine the eigenvalues and eigenvectors of these matrices.

Let $T_n(a, b, c) := \text{Tridiag}(\text{subdiag}, \text{diag}, \text{supdiag})$ denote the $n \times n$ tridiagonal matrix determined by $\text{subdiag} := (a, a, \dots, a, a)$, $\text{diag} := (b, b, \dots, b, b)$, and $\text{supdiag} := (c, c, \dots, c, c)$. In other words, $T_n(a, b, c)$ is the tridiagonal matrix that has all a 's on the subdiagonal, has all b 's on the diagonal and has all c 's on the superdiagonal. Recall that matrices with constant diagonals are called **Toeplitz** matrices. For example, when $n := 4$ we have:

$$T_4(a, b, c) := \begin{pmatrix} b & c & 0 & 0 \\ a & b & c & 0 \\ 0 & a & b & c \\ 0 & 0 & a & b \end{pmatrix}.$$

We want to find the eigenvalues and eigenvectors of the matrices $T_n(a, b, c)$.

The following result is well-known.

Proposition 3. *The matrix $T_n(a, b, c)$ is diagonally similar to the symmetric matrix $T_n(\sqrt{ac}, b, \sqrt{ac})$ provided $ac \neq 0$. In particular, $DT_n(a, b, c)D^{-1} = T_n(\sqrt{ac}, b, \sqrt{ac})$ where $D := \text{Diag}(1, d, d^2, \dots, d^n)$ and $d := \sqrt{c/a}$. Furthermore, $T_n(a, b, c)u = \lambda u$ iff $T_n(\sqrt{ac}, b, \sqrt{ac})Du = \lambda Du$ and hence the eigen-pairs of $T_n(a, b, c)$ are determined by the eigen-pairs of $T_n(\sqrt{ac}, b, \sqrt{ac})$.*

We call the transformation

$$T_n(a, b, c) \mapsto DT_n(a, b, c)D^{-1} = T_n(\sqrt{ac}, b, \sqrt{ac})$$

the **diagonal similarity symmetrizing transformation**.

Here is an example:

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & d & 0 \\ 0 & 0 & d^2 \end{pmatrix} \begin{pmatrix} b & c & 0 \\ a & b & c \\ 0 & a & b \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & d^{-1} & 0 \\ 0 & 0 & d^{-2} \end{pmatrix} \\ = \begin{pmatrix} b & c/d & 0 \\ da & b & c/d \\ 0 & da & b \end{pmatrix} = \begin{pmatrix} b & \sqrt{ac} & 0 \\ \sqrt{ac} & b & \sqrt{ac} \\ 0 & \sqrt{ac} & b \end{pmatrix}.$$

The following result is well-known. (See, for example, Rayleigh(1894).) We include a simple proof for the reader's convenience.

Proposition 4. Symmetric tridiagonal Toeplitz eigen-pairs. *The eigenvalues λ_j and corresponding eigenvectors u_j of the $n \times n$ matrix $T_n(a, b, a)$ are, for $j = 1, \dots, n$, given by $\lambda_j := b + 2a \cos(j\theta)$ and $u_j := (\sin(j\theta), \sin(2j\theta), \dots, \sin(nj\theta))^T$, where $\theta := \pi/(n+1)$.*

Proof. Recall that matrices A and $bI + A$ have the same eigenvectors. Also recall that λ is an eigenvalue of A iff $b + \lambda$ is an eigenvalue of $bI + A$.

Recall that, for $a \neq 0$, the matrices A and aA have the same eigenvectors. Also recall that λ is an eigenvalue of A iff $a\lambda$ is an eigenvalue of aA .

Using these reductions, we see that we only need to verify the eigen-pairs of $T_n(1, 0, 1)$. To reduce the number of symbols in this calculation we take $n = 3$. In this case, we have $\theta := \pi/4$ and

$$\begin{aligned} T_3(1, 0, 1)u_j &= \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} \sin(j\theta) \\ \sin(2j\theta) \\ \sin(3j\theta) \end{pmatrix} = \begin{pmatrix} \sin(0j\theta) + \sin(2j\theta) \\ \sin(1j\theta) + \sin(3j\theta) \\ \sin(2j\theta) + \sin(4j\theta) \end{pmatrix} \\ &= 2 \cos(j\theta) \begin{pmatrix} \sin(j\theta) \\ \sin(2j\theta) \\ \sin(3j\theta) \end{pmatrix}. \end{aligned}$$

The last equality follows from the well-known 3 term recurrence relation for the sine function:

$$\sin(k+1)\alpha - 2 \cos \alpha \sin(k\alpha) + \sin(k-1)\alpha = 0.$$

□

Proposition 5. Skew-symmetric tridiagonal Toeplitz eigen-pairs.

The eigenvalues λ_j and corresponding eigenvectors u_j of the $n \times n$ matrix $K_n := Z_n^T - Z_n$ are, for $j = 1, \dots, n$, given by $\lambda_j := 2i \cos(j\theta)$ and $u_j := (i \sin(j\theta), i^2 \sin(2j\theta), i^3 \sin(3j\theta), \dots, i^n \sin(nj\theta))$, where $\theta := \pi/(n+1)$.

Proof. Note that $K_n = Z_n^T - Z_n = T_n(-1, 0, 1)$. We apply the diagonal similarity symmetrizing transformation with $D^{-1} := \text{Diag}(1, i, i^2, \dots, i^n)$ to get

$$DT_n(-1, 0, 1)D^{-1} = T_n(i, 0, i) = iT_n(1, 0, 1).$$

We know the eigen-pairs of $T_n(1, 0, 1)$ from above. Thus we can easily determine the eigen-pairs of K_n . In particular, if $T_n(1, 0, 1)u = \lambda u$ then $DT_n(-1, 0, 1)D^{-1}u = iT_n(1, 0, 1)u = i\lambda u$. Hence $T_n(-1, 0, -1)(D^{-1}u) = i\lambda D^{-1}u$. $\square$

THE EIGENVALUES

Proposition 6. *The eigenvalues of the matrix R_n are 0 and $2i \cos(j\theta)$ for $j = 1, \dots, n-1$ where $\theta := \pi/n$.*

Proof. It is clear that the vector $(1, 1, \dots, 1)^T$ is an eigenvector with 0 as corresponding eigenvalue.

From the reduction proposition, we have

$$S_n^{-1}R_nS_n = K_n + e_n e_{n-1}^T.$$

Note that

$$K_n + e_n e_{n-1}^T = \begin{pmatrix} K_{n-1} & e_{n-1} \\ 0 & 0 \end{pmatrix}.$$

Also note that if $K_{n-1}u = \lambda u$ then

$$\begin{pmatrix} K_{n-1} & e_{n-1} \\ 0 & 0 \end{pmatrix} \begin{pmatrix} u \\ 0 \end{pmatrix} = K_{n-1}u = \lambda u = \lambda \begin{pmatrix} u \\ 0 \end{pmatrix}.$$

$\square$

Remark: The eigenvectors of R_n are determined in the proof of the proposition.

ACKNOWLEDGEMENTS

This paper was written in October of 2010 while Behn was visiting the mathematics department at Iowa State university (under grant number FONDECYT 1100135).

We thank Wayne Barrett (Brigham Young University) for his careful reading of this paper and his constructive comments about it.

Driessel thanks Wolfgang Kliemann, chair of the Mathematics Department at Iowa State, for arranging his affiliation with that department.

REFERENCES

- Drew, J.H.; Johnson, C.R.; Olesky, D.D.; and van den Driesche, P. (2000) Spectrally arbitrary patterns, *Linear Algebra and Appl.* 308, 121-137
- Elsner, L.; Olesky, D.D.; and van den Driesche, P. (2003) Low rank perturbations and the spectrum of a tri-diagonal sign pattern, *Linear Algebra and Appl.* 308, 121-137
- Kailath, T.; Kung, S.Y.; and Morf, M. (1979a) Displacement ranks of matrices and linear equations, *J. Math. Anal. Appl.* 68, 395-407
- Kailath, T.; Kung, S.Y.; and Morf, M. (1979b) Displacement ranks of a matrix, *Bull. Amer. Math. Soc.* 1, 769-773
- Kailath, T. and Sayed, A.H. (1999) *Fast Reliable Algorithms for Matrices with Structure*, SIAM
- Rayleigh, J.W.S. (1894) *The Theory of Sound*, Macmillan (Reprinted by Dover in 1945.)
- Trench, W.F. (2004) Characterization and properties of matrices with generalized symmetry or skew symmetry, *Lin. Alg. Appl.* 377, 207-218
- Weaver, J.R. (1985) Centrosymmetric (Cross-Symmetric) Matrices, *American Mathematical Monthly* 92, 711-717